arXiv:1603.07489v1 [math.NT] 24 Mar 2016

Modular Curves with many Rational Points

Valerio Dose, Pietro Mercuri, Claudio Stirpe

Abstract

Using classical and non split modular curves we improve many lower bounds for the maximum number of points in a curve over finite fields.

1 Introduction

In recent papers, Pietro Mercuri [1], [2] found explicit equations defining the modular curves $X_{ns}^{+}(p)$ for $p = 17$, 19 and 23 and $X_0^{+}(p)$ for $p = 163$, 197, 229, 269 and 359, where $X_{ns}(p)$ is the modular curve associated to the normalizer of a nonsplit Cartan subgroup of the general linear group over a finite field with p elements and $X_{ns}^{+}(p)$ is the quotient by the Atkin involution. Very few is still known about $X_{ns}^{+}(p)$ and they are harder to study with respect to modular curves $X_0^{+}(p)$ considered here. Such curves are also interesting from the point of view of the number of points. In this paper we use explicit equations to improve the upper bound for the maximum number of places in algebraic curves over finite fields. Modular curves, in fact, provide many interesting examples which could improve the results listed, at time of writing (March 2016), in the website manypoints.org, see [3]. We could improve some lower bounds for curves of genus $g = 5$, 7, 8 and 13.

2 Basic Facts

We recall that the Weil bound prescribes the following inequality

$$N \leq q + 1 + 2g\sqrt{q},$$

where N is the number of rational points of a smooth, projective, absolutely irreducible curve of genus g over the finite field $\mathbb{F}_q$ with q a power of a prime number ℓ. The previous estimate may be sharp only for small values of g. Since 1980 many authors provided sharper estimates for several choices of g and q. The reader may check the full list of the upper and lower bounds for the maximum value of N for $g \leq 50$ in the table in [3]. In this section we improve some lower bounds for $g = 5$, 7, 8 and 13. As in [3], when the lower bound is missing we consider a curve interesting if $M_g/N < \sqrt{2}$, where M_g is the upper bound for the number of rational point for a curve of genus g over the finite field $\mathbb{F}_q$.

3 Lower bounds for the maximum number of points

In this section we list explicit examples of curves with many rational points over finite fields. We start from the case $g = 5$.

3.1 Improved lower bound for $g = 5$

Let C be the classical modular curve $X_0(73)$. Explicit equations are already available on MAGMA. The reader may check using a computer algorithm that such curve has genus 5. The reduction modulo ℓ gives many interesting results when $\ell = 5, 11$ and 13.

We start with the case $\ell = 5$. The reduction of C modulo 5 is good because $5 \nmid 73$. We compute using MAGMA that the curve C has 58 rational points over $\mathbb{F}_{5^2}$ and 642 rational points over $\mathbb{F}_{5^4}$. Now let $\ell = 11$. Since $11 \nmid 73$ also the reduction modulo 11 is good and the number of $\mathbb{F}_{11^3}$-rational points for C is 1270. Finally, for $\ell = 13$, the reduction modulo 13 is again good and gives two interesting results. In fact, C has 234 rational points over $\mathbb{F}_{13^2}$ and 2156 rational points over $\mathbb{F}_{13^3}$. At the moment, such values are the best known, according to [3]. We summarize the previous results in the following table.

Curve	q	N	M_5
$X_0(73)$	5^2	58	72
$X_0(73)$	5^4	642	876
$X_0(73)$	11^3	1270	1692
$X_0(73)$	13^2	234	300
$X_0(73)$	13^3	2156	2663

3.2 Improved lower bound for $g = 6$

Let C be the genus 6 curve $X_0^+(271)$. Then C has good reduction modulo $\ell = 3$. The number of rational points over $\mathbb{F}_{3^2}$ is 35. According to the tables in [3] this value attaines the lower bound for the maximum number of rational points for a genus 6 curve.

3.3 Improved lower bound for $g = 7$

Consider the following genus 7 curves $X_0^+(193), X_0^+(229), X_0^+(257)$ and $X_0(281)^+$. All these curves have good reduction modulo $\ell = 5, 7, 11, 13, 31$. In the following table we list the most interesting values.

Curve	q	N	M_7
$X_0^+(193)$	5^4	808	976
$X_0^+(193)$	11^3	1357	1836
$X_0^+(229)$	5^5	3242	3903
$X_0^+(257)$	13^3	2156	2663
$X_0^+(281)$	7^5	17260	18617
$X_0^+(281)$	13^2	271	352
$X_0^+(281)$	31	73	103

3.4 Improved lower bound for $g = 8$

The genus 8 curve $C = X_0^+(293)$ has good reduction modulo $\ell = 5, 7, 11, 13, 17, 19$. Similarly we can reduce $X_{ns}^+(19)$ by any prime $l \neq 19$. Some interesting values are listed in the table.

Curve	q	N	M_8
$X_0^+(293)$	5^5	3177	4014
$X_0^+(293)$	7^5	16736	18875
$X_0^+(293)$	11^4	14863	16579
$X_0^+(293)$	19^3	6824	8180
$X_{ns}^+(19)$	13^2	316	378
$X_{ns}^+(19)$	13^3	2309	2942
$X_{ns}^+(19)$	17^3	5247	6032

Remark 3.1. We notice that both curves have many points on the same fields $\mathbb{F}_q$. As an Example $X_0^+(293)$ over $\mathbb{F}_{13^3}$ has 2101 points. This is less than the value in the previous table but it is also an interesting result. Similar results hold for $q = 13^2$ and $q = 17^3$. Similarly $X_{ns}^+(19)$ has many points over $\mathbb{F}_q$ for $q = 11^4$, 5^5 and 7^5. It is a very interesting question to see if there is a common lower bound for the number of points over $\mathbb{F}_q$ for other values of q.

3.5 Improved lower bound for $g = 13$

Finally $X_{ns}^+(23)$ has genus 13. As in the previous case we reduce for a prime l not dividing 23. Here we list an other interesting result.

Curve	q	N	M_8
$X_{ns}^+(23)$	17^2	518	732

References

[1] P. Mercuri, *Equations of Modular Curves Associated to Normalizers of non-split Cartan Subgroups*, preprint 2016.

[2] P. Mercuri, *Equations and Rational Points of the Modular Curves* $X_0^+(p)$, preprint 2016.

[3] G. van der Geer, E. Howe, K. Lauter, C. Ritzenthaler, *Tables of Curves with Many Points*, (2009), http://www.manypoints.org.